

Power-law behavior in signal scattering process in vertical granular chain with light impurities

Aiguo Xu

Department of Physics and
Center for Strongly Correlated Materials Research,
Seoul National University, Seoul 151-742, Korea

February 2, 2008

Abstract

We investigate the scattering process of impulse in vertical granular chain with light impurities. When the perturbation is weak, the quantities describing the reflection rate exhibit power law behavior with the impurity depth. The exponent is nearly independent of v_i . When the perturbation is very strong, the vertical chain shows similar behavior to that of the horizontal chain, so the exponent is zero. Our numerical investigation begins from the weak perturbation region and extend to the nonlinear region and found a peak of the exponent. The difficulty in extending the numerical investigation to a stronger perturbation region is analyzed.

PACS numbers: 45.70-n, 43.25.+y, 46.40.Cd

Key Words: granular chain, scattering process, power-law behavior, reflection rate

1 Introduction

In the last ten years physicists have been interested again in the study of dry granular materials. Granular materials are ubiquitous around us. They have many properties that are greatly different from those associated with common

solids, liquids and gases. These unique properties lead to many applications. Many new ideas have emerged from this recent development. These systems have been studied from various different viewpoints experimentally, numerically, and theoretically[1, 2, 3]. Nesterenko[4]noted that the propagation of a perturbation in loaded chain with Hertzian contacts possesses soliton-like features. The soliton-like behavior of the signal in two-dimensional granular beds was studied by Sinkovits and Sen[5, 6]. The soliton-like properties of the signal in the chain of iron balls were examined experimentally by Coste et al[7]. Sen et al [8] studied the soliton-like pulse in perturbed and driven Hertzian chains and their possible applications in detecting buried impurities. For the vertical granular chain under gravity with power-law-type contact force, J. Hong et al [9] showed that there are two kinds of propagating modes, quasisolitary and oscillatory, depending on the strength of impulse. The type of dispersion and the oscillation frequency, wavelength, and period follow power laws in depth; J. Hong and Aiguo Xu[10] found that the signal speed follows a power law in depth and the exponent varies smoothly from a saturant value to zero when the strength of the perturbation varies from very weak to very strong.

To explore the prospects for the application of impulse for detecting certain buried objects in granular beds, it is imperative that one must acquire a detailed understanding of the its properties. Up to now many problems on this topics are still openning. In present paper, we study the scattering process of impulse when it encounters some other kind of grains which we name impurities. The granular chain system can be described by the following motion equations

$$\begin{aligned} m_n \ddot{x}_n &= V'(x_{n-1}, x_n) - V'(x_n, x_{n+1}) + m_n g, \\ n &= 1, 2, \dots, N. \end{aligned} \tag{1}$$

where m_n is the mass of the n th grain, $V(x_n, x_{n+1})$ is the inter-grain interaction between the n th and the $(n + 1)$ th grains. In this paper we study the impulses in a special case with $V(x_{n-1}, x_n) = [\eta/(p+1)][\Delta_0 - (x_n - x_{n-1})]^{p+1}$, where Δ_0 is the diameter of the spherical grain, p the exponent of the power-law type contact force, and η the elastic constant of the grain under consideration. Thus equation (1) becomes

$$\begin{aligned}
m_n \ddot{x}_n &= \eta \{ [\Delta_0 - (x_n - x_{n-1})]^p - [\Delta_0 - (x_{n+1} - x_n)]^p \} \\
+m_n g, n &= 1, 2, \dots, N.
\end{aligned} \tag{2}$$

For the Hertzian chain, i.e., the chain with Hertzian interaction between neighboring granular spheres, $p = 3/2$.

2 Analytic treatment

In our system most of the grains are same and the second kind of grains only occur as impurities. Hence to analytically get some information of the system, we first fix the mass of all the grains to a constant, i.e., $m_n = m$ for all n . Then we introduce a new variable ψ_n , denoting the displacement of n th grain from equilibrium, defined by

$$\psi_n = x_n - n\Delta_0 + \sum_{l=1}^n \left(\frac{mgl}{\eta} \right)^{1/p}, \tag{3}$$

where the last term in the right-hand side is the sum of grain overlaps up to n th contact and we set $x_0 = \psi_0 = 0$. Eq. (2) is transformed into

$$\begin{aligned}
m \ddot{\psi}_n &= \eta \left[\left(\frac{mgn}{\eta} \right)^{1/p} + (\psi_{n-1} - \psi_n) \right]^p \\
&- \eta \left[\left\{ \frac{mg(n+1)}{\eta} \right\}^{1/p} + (\psi_n - \psi_{n+1}) \right]^p + mg
\end{aligned} \tag{4}$$

using Eq. (3).

For the weakly nonlinear regime, the condition

$$|\psi_{n-1} - \psi_n| \ll \left(\frac{mgn}{\eta} \right)^{1/p} \tag{5}$$

is valid and the expansion of Eq. (4) under this condition reads

$$m \ddot{\psi}_n = -\mu_n (\psi_n - \psi_{n-1}) + \mu_{n+1} (\psi_{n+1} - \psi_n) \tag{6}$$

where $\mu_n = mpg \left(\frac{\eta}{mg} \right)^{1/p} n^{1-\frac{1}{p}}$ is the force constant of n th contact of the linear horizontal chain. We drop weak nonlinear terms of the expansion in Eq. (6) to make the system linear. Thus the vertical granular chain becomes a horizontal chain with varying force constants in which the gravity effect is contained. Both left and right side of Eq. (6) are linear in ψ_n . Therefore, the scaling analysis tells us that the equation of motion (6) has nothing to do with the initial impulse v_i . For this case we [10] have obtained the following power-law,

$$v(h) \propto h^{-\frac{1}{4}(\frac{1}{3}+\frac{1}{p})}. \quad (7)$$

The reducing of signal amplitude mainly due to the dispersion effect.

In our present studies two distances are important. The first one is the depth of the impurities d_{imp} . The second one is the distance between the impurity center and the grain where we get datas, d . Our main objectives are to get some knowledge on the reflection rate and transmission rate of the kinetic energy from the impurities. For fixed d , we expect the reflection rate follows a power-law and the exponent is independent of v_i .

For the strongly nonlinear regime, the condition

$$(\psi_{n-1} - \psi_n) \gg \left(\frac{mgn}{\eta} \right)^{1/p} \quad (8)$$

is valid and the expansion under this condition leads Eq. (4) to

$$\begin{aligned} m\ddot{\psi}_n &= \eta [(\psi_{n-1} - \psi_n)^p + pg_n(\psi_{n-1} - \psi_n)^{p-1}] \\ &- \eta [(\psi_n - \psi_{n+1})^p + pg_{n+1}(\psi_n - \psi_{n+1})^{p-1}] \end{aligned} \quad (9)$$

where $g_n = (mgn/\eta)^{1/p}$ denotes grain overlap at n th contact. The gravity term can be neglected in the highly nonlinear regime, since the gravity effect appears in the coefficient g_n . Different order of ψ_n in the left and right side of Eq. (9) implies that v_i -dependence must be appeared in the signal characteristics. Under the limitation $v_i \rightarrow \infty$, Eq. (9) recovers to

$$m\ddot{\psi}_n = \eta[(\psi_{n-1} - \psi_n)^p - (\psi_n - \psi_{n+1})^p] \quad (10)$$

That is to say, the vertical chain shows same behavior as that of the horizontal chain. For the horizontal chain the signal speed is a constant, so the exponent of the depth-dependence is 0.

3 Simulation results

Our results are based on careful numerical integration of the coupled equations of motion for granular chains with N spheres, where N varies from 1000 to 5000 according to our need. The predictor-corrector method, Runge-Kutta method, Richardson extrapolation and the Bulirsch-Stoer method[11] are used. The program units are just the same as those in Refs.[8] and [9]. That is to say, the units of distance, mass, and time are $10^{-5}m$, $2.36 \times 10^{-5}kg$, and $1.0102 \times 10^{-3}s$, respectively. This kind of program units gives the gravitational acceleration $g = 1$. We set the grain diameter 100 and $\eta = 5657$.

Figure 1 shows four snapshots for a scattering process, where the impurity depth is $d_{imp} = 500$, which is described by the grain number; there are five impurities around the depth $d_{imp} = 500$; the mass of the impurity grain is $m_{imp} = 0.1$; the initial perturbation strength is $v_i = 0.1$. Fig. 1 (a) shows a snapshot before the scattering process. Figs. 1 (b) and 1(c) show the scattering process. Fig. 1 (d) shows a snapshot where the scattering process nearly finished. We use “1st” and “2nd” to denote the leading and second leading peaks. The leading and second leading peaks of the incident impulse are pointed by an arrow, respectively, in Figs. 1 (a) and 1(b). The leading and second leading peaks of the scattered impulse are pointed by an arrow, respectively, in Figs. 1 (c) and 1 (d). It is clear that the leading peak of the scattering impulse is upwards, which is a characteristic property of the scattering process from light impurities[6]. We use five impurities because that the scattered impulse is too weak if we use only one impurity. To understand the scattering process, we are interested in the variation of the ratios $v_{ref}^{1st}/v_{inc}^{1st}$, $v_{ref}^{2nd}/v_{inc}^{2nd}$ and $v_{ref}^{2nd}/v_{inc}^{1st}$ with the depth of the impurities d_{imp} , where v_{inc}^{1st} and v_{ref}^{1st} are maximum values of the leading peaks of the incident and reflected impulses, v_{inc}^{2nd} and v_{ref}^{2nd} are maximum values of the second leading peaks of the incident and reflected impulses. We obtain these datas from a gain with depth $n = d_{imp} - d$. The three ratios describe the reflection rate from different sides. During the scattering process the shape of the impulse is not stable, so an appropriate distance d is necessary. These ratios describe the reflection rate of the kinetic energy in the vertical granular chain. In order to get exact value for the leading and second leading peaks, we use the smooth velocity-time curve of the measured gain, instead of the rough snapshots.

Fig.2 shows the variations of $v_{ref}^{1st}/v_{inc}^{1st}$, $v_{ref}^{2nd}/v_{inc}^{2nd}$ and $v_{ref}^{2nd}/v_{inc}^{1st}$ with the

impurity depth d_{imp} . Fig. 2 (a) is for $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ and $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$, where $d = 200$, $v_i = 1$. The squares are numerical results for $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ and the circles are numerical results for $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$. The lines are corresponding fitting curves. Fig.2 (b) is for $v_{ref}^{2nd}/v_{inc}^{1st}(d_{imp})$, where $d = 200$, $v_i = 5$. The solid squares are for numerical results and the line is the fitting curve. We use the simplest power-law model, $v_{ref}/v_{inc} \propto d_{imp}^{-\beta}$, to fit the numerical results. The fitting results suggest that the three ratios follow power-laws when d_{imp} is large enough. From Refs. [9] we know that when the initial perturbation is weak the propagating mode is oscillatory, when the initial perturbation is strong the propagating mode is quasisolitary. It is interesting to check if the power-law model works for both the weak and strong perturbations. If it works, we have the following expected result: When v_i is large enough, the behavior of the vertical chain is similar to that of the horizontal chain; hence there is no power-law. That means $\beta \rightarrow 0$ as $v_i \rightarrow \infty$.

Fig.3 shows the variation of the exponent for $v_{ref}^{2nd}/v_{inc}^{2nd}$ with the strength of initial perturbation, where the line with squares is for the case of $d = 100$, the line with circles is for the case of $d = 200$. The two curves show similar behaviors. It is clear that the exponent is independent of v_i when v_i is very small. For the case of $d = 100$, $\beta = 0.19 \pm 0.01$. For the case of $d = 200$, $\beta = 0.23 \pm 0.01$. In fact when v_i is too small we can hardly get helpful information from the simulation datas due to numerical errors. The curve for $d = 100$ has a peak around $\tilde{v}_i = 5$. The curve for $d = 200$ has a peak around $\tilde{v}_i = 10$. When $v_i > \tilde{v}$, β decreases quickly with the increasing of v_i . We tried to add more points to the figure, but when v_i is larger than 20 we can hardly measure the second leading peak from the incident impulse. We can know the reason from Fig. 4.

Fig. 4 shows a $v_n(t)$ relation, where $d_{imp} = 300$, $d = 200$, $n = 100$, $v_i = 15$, and $m_{imp} = 0.1$. The impulse near $t = 2$ is the incident one. The impulse within $9 < t < 10$ is the reflected one (Only part of the reflected impulse is plotted). It is clear that for a strong perturbation the propagating impulse is quasisolitary and it has no evident the second leading peak; at the same time, the leading peak of the reflected impulse is very small and there exist some oscillations before the leading peak occurs.

When the perturbation is weak, the propagation of the wave is oscillatory; hence the two ratios $v_{ref}^{1st}/v_{inc}^{1st}$ and $v_{ref}^{2nd}/v_{inc}^{2nd}$ can describe well the reflection rate of the kinetic energy from the impurities. But with the increasing of

v_i , the propagation of the wave is more and more quasisolitary and the ratio $v_{ref}^{2nd}/v_{inc}^{1st}$ describes the reflection rate better. Fig.5 shows the variation of the exponent β' for $v_{ref}^{2nd}/v_{inc}^{1st}(d_{imp})$ with v_i , where $d = 200$. It shows similar behavior to the exponent β for $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$. When v_i is very small, $\beta' \approx 0.2$.

Although we have found the power-law behavior for $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ for not very large v_i (see Fig. 2 for an example), it is hard for us to get a figure which is similar to Fig.3 and Fig.5. The reasons will be given in the following part.

For a fixed distance d , when the depth of impurities d_{imp} is large, we find the curve $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ oscillates greatly with d_{imp} . Fig. 6 shows two examples, where $d = 200$; $v_i = 6$ in (a) and $v_i = 4$ in (b). We can understand the reason for this kind of behaviors through Fig. 7. Fig.7 shows several $v_n(t)$ curves from which we can study the scattering process of the impulse, where $n = d_{imp} - 100$, $v_i = 0.1$, $m_{imp} = 0.1$; $d_{imp} = 600$ in (a), $d_{imp} = 800$ in (b), $d_{imp} = 1000$ in (c), and $d_{imp} = 1100$ in (d), respectively. For the reflected impulse, only the leading peak and the second peak are shown. For a fixed value of d , when the depth d_{imp} is small, the reflected impulse and the incident impulse are separated, i.e. there is a time during which the measured grain is static. For these cases, we can get exact values for v_{inc}^{1st} , v_{ref}^{1st} , v_{inc}^{2nd} , and v_{ref}^{2nd} . When the depth d_{imp} is large enough, the tail of the incident impulse and the reflected impulse are connected. For these cases, we can hardly get exact values for v_{ref}^{1st} and v_{ref}^{2nd} . From Fig.7 we also understand the fact these phenomena first influence the measurement of v_{ref}^{1st} , then v_{ref}^{2nd} .

We name the increasing process of d_{imp} before a power-law occurs a transient process. The simulation results show that the ratio $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ needs a much longer transient process than $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$. The initial impulse is stronger, we need a longer transient process. The curves for $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$ decreases with the increasing of d_{imp} , while the curves for $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ first increases, then decreases. The initial impulse is stronger, the curves decrease more slowly. We also notice that there are more fitting errors when v_i is large.

4 Conclusion

In present paper we study the scattering process of impulse in vertical granular chain with light impurities. Most of the studies resort on numerical simulations and the analytic treatment helps to understand the behaviors. When the perturbation is weak, the quantities describing the reflection rate exhibit power law behavior with the impurity depth. The exponent is independent of v_i . When the perturbation is very strong, the vertical chain shows similar behavior to that of the horizontal chain, so the exponent is zero. Our numerical investigation begins from the weak perturbation region and extend to the nonlinear region and found a peak of the exponent. The difficulty in extending the numerical investigation to a stronger perturbation region is analyzed. More work needs to be done to completely understand the scattering process of the signal in vertical granular chain.

Acknowledgments

The author acknowledges Prof. Jongbae Hong for helpful suggestions and discussions. This work was supported by Korea Research Foundation Grant (KRF-2000-DP0106). The author also wishes to acknowledge the support by the BK21 project of the Ministry of Education, Korea.

References

- [1] *Physics of Dry Granular Media*, edited by H.J.Herrmann, J-P.Hovi, and S.Luding (Kluwer Academic Publishers, Dordrecht, 1998)
- [2] *Powders and Geains 97: Proceedings of the 3rd International Conference on Powders and Grains, North Carolina, 1997*, edited by R.P.Behringer and J.T.Jenkins (A.A.Balkema, Rotterdam, 1997).
- [3] H.M.Jaeger and S.R.Nagel, Science 255, 1523 (1995); H.M.Jaeger, S.R.Nagel, and R.P.Behringer, Rev. Mod. Phys. **68**, 1259 (1996).
- [4] V.F.Nesterenko, J.Appl. Mech. Tech. Phys. 5 733 (1984); J.Phys. IV 55, C8-729 (1994).
- [5] R.S.Sinkovits and S.Sen, Phys. Rev. Lett. 7, 2686 (1995).
- [6] S.Sen and R.S.Sinkovits, Phys. Rev. E 5, 6857 (1996).

- [7] C.Coste, E.Falcon, and S.Fauve, Phys. Rev. E 56, 610 (1997).
- [8] S.Sen, M.Manciu, and J.D.Wright, Phys. Rev. E 57, 2386 (1998).
- [9] J. Hong, J.Y. Ji, and H.Kim, Phys. Rev. Lett. 82, 3058 (1999); J. Hong, H. Kim, and J. P. Hwang, Phys. Rev. E 61, 964 (2000).
- [10] J. Hong and Aiguo Xu, Phys. Rev. E **63**, 061310 (2001).
- [11] *Numerical Recipes in FORTRAN, The Art of Scientific Computing*, second edition, edited by W. H. Press, S. A. Teulolsky, W. T. Vetterling, B. P. Flannery (Cambridge University Press, 1992).

Figure caption:

Figure 1: Snapshots for a scattering process, where the impurity depth is $d_{imp} = 500$, which is described by the grain number; there are five impurities around the depth $d_{imp} = 500$; the mass of the impurity grain is $m_{imp} = 0.1$; the initial perturbation strength is $v_i = 0.1$.

Figure 2: Variations of $v_{ref}^{1st}/v_{inc}^{1st}$, $v_{ref}^{2nd}/v_{inc}^{2nd}$ and $v_{ref}^{2nd}/v_{inc}^{1st}$ with the depth of the impurities d_{imp} . Fig.2 (a) is for $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ and $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$ where $d = 200$, $v_i = 1$. The squares are numerical results for $v_{ref}^{1st}/v_{inc}^{1st}(d_{imp})$ and the circles are numerical results for $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$. The lines are corresponding fitting curves. Fig.2 (b) is for $v_{ref}^{2nd}/v_{inc}^{1st}(d_{imp})$ where $d = 200$, $v_i = 5$. The solid squares are for numerical results and the line is the fitting curve.

Figure 3: Variation of the exponent for $v_{ref}^{2nd}/v_{inc}^{2nd}$ with the strength of initial perturbation, where the line with squares is for the case of $d = 100$, the line with circle is for the case of $d = 200$.

Figure 4: The $v_n(t)$ relation, where $d_{imp} = 300$, $d = 200$, $n = 100$, $v_i = 15$, and $m_{imp} = 0.1$. The impulse near $t = 2$ is the incident one. The impulse within $9 < t < 10$ is the reflected one (Only part of the reflected impulse is plotted). The second leading peak of the incident impulse and the leading peak of the reflected impulse are very small. Some oscillations occur behind the main incident impulse.

Figure 5: Variation of the exponent β' for $v_{ref}^{2nd}/v_{inc}^{1st}(d_{imp})$ with v_i , where $d = 200$. It shows similar behavior to the exponent β for $v_{ref}^{2nd}/v_{inc}^{2nd}(d_{imp})$. When v_i is very small, $\beta' \approx 0.2$.

Figure 6: Reflection rate $v_{ref}^{1st}/v_{inc}^{1st}$ versus d_{imp} , where $d = 200$; $v_i = 6$ in (a) and $v_i = 4$ in (b).

Figure 7: Velocity $v_n(t)$ curves, where $n = d_{imp} - 100$, $v_i = 0.1$, $m_{imp} = 0.1$; $d_{imp} = 600$ in (a), $d_{imp} = 800$ in (b), $d_{imp} = 1000$ in (c), and $d_{imp} = 1100$ in (d), respectively.















